

DANGER
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



Coxidin 200

Active Ingredient:

Contains monensin sodium, equivalent to 200 g/kg MONENSIN activity

Net contents: 20 kg

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

INDICATIONS:

Dairy Cattle:

For increased milk protein production. As an aid in the control of ketosis. As an aid in the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

Beef Cattle:

For improved feed efficiency and/or rate of weight gain. As an aid in the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

Beef Cows:

For improved feed efficiency as indicated by improvement in the relationship of feed to cow weight, calf weight and reproductive performance.

Replacement Heifers:

For increased rate of weight gain and to promote earlier onset of first oestrous. As an aid in the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

Sheep:

As an aid in the control of coccidiosis caused by *Eimeria ninakholyakimovae*, *E. ahsata*, *E. faurei*, *E. parva*, *E. intricata* and *E. pallida*.

Goats:

As an aid in the control of coccidiosis caused by *Eimeria arloingi*.

Broiler Chickens and Layer Replacement Chickens:

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*.

Growing Turkeys:

As an aid in the prevention of coccidiosis caused by *Eimeria adenoides*, *E. gallopavonis* and *E. meleagrimitis*.

MIXING AND FEEDING DIRECTIONS

IMPORTANT:

MUST BE THOROUGHLY MIXED IN FEEDS BEFORE USE.

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Feedlot cattle: In complete feeds:

Thoroughly mix COXIDIN 200 into the ration at a rate of 50 g to 165 g COXIDIN 200 per tonne of final feed. This is equivalent to 10 to 33 ppm (mg/kg) of monensin active ingredient in the final feed.

In Supplements:

Thoroughly mix COXIDIN 200 into the supplement to provide the equivalent of 50-360 mg monensin per animal per day, when the supplement is fed at its recommended rate. This is equivalent to 0.25 - 1.8 g Coxidin 200 per animal per day.

Calves : In Supplements:

Thoroughly mix COXIDIN 200 into the supplement to provide the equivalent of 100 ppm (mg/kg) of monensin active ingredient in the supplement. This is equivalent to administering 0.5g of COXIDIN 200 per calf / day when supplement is fed at 1kg / calf / day.

Beef cattle, Beef cows, Replacement Heifers:

In Supplements:

Thoroughly mix COXIDIN 200 into the supplement to provide 200 mg monensin per animal per day when the supplement is fed at its recommended rate. This is equivalent to feeding 1g of COXIDIN 200 per animal per day.

Daily Supplement Provided (kg/hd/day)	0.5	1.0	2.0
COXIDIN 200 Addition Rate (kg/t)	2	1	0.5

Dairy cows:

In Supplements:

Thoroughly mix COXIDIN 200 into the supplement to provide 300 mg monensin per animal per day when the supplement is fed at its recommended rate. This is equivalent to feeding 1.5g of COXIDIN 200 per animal per day. Do not use unless provided more than 0.5kg feed.

All rations (including dry feeds, silage, wet feeds) should be converted to a 90% dry matter basis for calculation of the COXIDIN 200 addition rate.

Daily Supplement Provided (kg/hd/day)	1.0	2.0	3.0
COXIDIN 200 Addition Rate (kg/t)	1.5	0.75	0.5

Sheep and goats:

Complete feed:

Thoroughly mix COXIDIN 200 into the ration at a rate of 25 to 100 g per tonne of final feed. This is equivalent of 5 to 20 ppm (mg/kg) of monensin active ingredient in the final feed.

In Supplements:

Thoroughly mix COXIDIN 200 into the supplement to provide 7.5 mg to 40 mg monensin per animal per day when the supplement is fed at its recommended rate. This is equivalent to feeding 0.0375 - 0.2 g of COXIDIN 200 per animal per day.

Example: In order to provide 30 mg / hd / day monensin intake:

Daily Supplement Provided (kg/hd/day)	0.1	0.2	0.3
COXIDIN 200 Addition Rate (kg/t)	1.5	0.75	0.5

Broiler and Layer Replacement Chickens:

Thoroughly mix 0.5 kg to 0.6 kg of Coxidin 200 in one tonne of feed to provide monensin equivalent to 100g to 120g activity per tonne of complete feed.

Growing Turkeys:

Thoroughly mix 0.3 kg to 0.5 kg of Coxidin 200 in one tonne of feed to provide monensin equivalent to 60g to 100g activity per tonne of complete feed.

Feeding Directions:

Broiler Chickens:

Feed continuously as the only ration.

Layer Replacement Chickens:

May be fed up to the point of laying.

Growing Turkeys:

Feed continuously as the only ration until 16 weeks of age.

CONTRAINDICATIONS:

Poultry consuming monensin should not be treated with products containing tiamulin or oleandomycin.

Severe growth depression may occur.

CAUTION:

Do not allow use on horses or dogs as fatal toxicosis may result. Ensure recommended doses are not exceeded. Care must be exercised when feeding concurrently with other antimicrobials.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

Cattle:

Meat: Nil

Milk: Nil

Sheep and Goats:

Meat: Nil

Milk: Nil

Milk intended for human consumption must be discarded during treatment for not less than **35 days** following the last treatment.

Poultry:

Meat: Nil.

Eggs from treated birds must not be sold for human consumption for

10 days following the last treatment.

DANGER:

Corrosive

Dangerous to the environment

Handling precautions:

Monensin sodium may be dangerous if swallowed or absorbed through the skin and may cause eye damage.

Repeated exposure to monensin sodium may cause skin allergy. May cause mild skin irritation. Avoid skin and eye contact. When mixing wear impervious gloves. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face before meals and after work.

First aid:

If swallowed do NOT induce vomiting. If eye contact occurs hold eyes open and rinse cautiously with water for several minutes. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

Environmental Protection:

Harmful to aquatic organisms and soil environment. Very toxic to terrestrial vertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

Emergency Response:

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material.

Disposal:

Preferably dispose of product by use, or at an approved landfill or equivalent facility.

STORAGE:

Store below 25°C. Protect from direct sunlight. Store in its original container in a dry, well-ventilated area.

Registered pursuant to the ACVM Act 1997, No. A10328

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Registered to: AgriHealth NZ

www.agrihealth.co.nz

Manufactured by:

Huvepharma AD 33 James Boucher Blvd.

1407 Sofia, Bulgaria

www.huvepharma.com

Distributed by: AgriHealth NZ

Phone 09 889 8210

www.agrihealth.co.nz

Batch:0000000000

Expiry:00.2000

LB530v2Cox200mp20-NZ0111