



FOR ANIMAL TREATMENT ONLY  
RESTRICTED VETERINARY MEDICINE  
**Pharmasin<sup>®</sup> 25%**  
**Granular premix**

**TYLOSIN PHOSPHATE**  
**FOR USE IN CHICKENS, BEEF CATTLE and PIGS**

**Active Ingredient:** TYLOSIN (as TYLOSIN PHOSPHATE) 250 g/kg

Tylosin contains not less than 80% factor A

**Net contents: 20 kg**

**READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT**

**INDICATIONS:**

**Beef Cattle:**

For the reduction in the incidence of liver abscess.

**Broiler Chickens and Replacement Layers:**

As an aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum* and/or *Mycoplasma synoviae*. For the treatment and prevention of necrotic enteritis associated with susceptible strains of *Clostridium perfringens*.

**Pigs:**

For the treatment and prevention of Ileitis.

**MIXING AND FEEDING DIRECTIONS:**

**IMPORTANT: MUST BE THOROUGHLY MIXED IN FEEDS BEFORE USE.**



Animal Species	Recommended use	Feeding period	Weight of Pharmasin 25% Granular Premix per 1000kg of Complete Feed	Use Level
<b>Pigs</b>	For the treatment and prevention of Ileitis. Prevention-length of treatment required will be depend on the level of disease. Consult your vet for advice.	Treatment prevention	400 g 160 g	100 ppm 40 ppm
<b>Broiler Chickens and Replacement Layers</b>	As an aid in the control of chronic respiratory disease caused by <i>Mycoplasma gallisepticum</i> and/or <i>Mycoplasma synoviae</i> . For the treatment and prevention of necrotic enteritis associated with susceptible strains of <i>Clostridium perfringens</i> .	For 5 days after hatching. Repeat for 2 days at 3 to 5 weeks of age.  Treatment-treat for 7 days. Prevention-feed during periods of risk of clinical disease	3600 g to 4000 g  400 g 400 g	900 ppm to 1000 ppm  100 ppm 100 ppm
<b>Layers</b>	As an aid in the control of chronic respiratory disease caused by <i>Mycoplasma gallisepticum</i>	Feed from time of shifting into laying cages	88 g to 220 g	22 ppm to 55 ppm
<b>Beef Cattle</b>	For reduction in the incidence of liver abscess. For supplementary feeds 60-90 mg tylosin/head/day	Feed during period on ration	44 g	11 ppm

The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. Indiscriminate use of this product could contribute to antibiotic resistance. The product should only be used after consideration of the alternative therapeutic and preventative measures.

**WITHHOLDING PERIODS:**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.**

Meat: Cattle Nil  
Pigs Nil  
Poultry Nil  
Eggs: Nil

**WARNING**

**Dangerous to the Environment**

**Handling precautions:**

Tylosin may cause respiratory sensitization. Repeated exposure to tylosin may cause skin allergy. Avoid inhalation and contact with the skin.

KEEP OUT OF REACH OF CHILDREN

**Environmental Protection:**

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container.

**Disposal:**

Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility

**STORAGE:**

Store in original pack, in a dry place, below 25°C. Protect from light. Do not refrigerate or freeze - protect from frost.

Restricted Veterinary Medicine. Registered pursuant to the ACVM Act 1997, No. A010451  
See [www.foodsafety.govt.nz/industry/acvm/](http://www.foodsafety.govt.nz/industry/acvm/) for registration conditions.

Approved under the HSNO Act, No. HSR002370. Refer [www.ermanz.govt.nz](http://www.ermanz.govt.nz) for controls

**Manufactured by:**

Huvepharma AD  
[www.huvepharma.com](http://www.huvepharma.com)

**Distributed by:**

AgriHealth NZ Ltd  
[www.agrihealth.co.nz](http://www.agrihealth.co.nz)

Batch:

Expiry:

**AgriHealth**



GRANULAR PREMIX

CHICKENS, BEEF CATTLE AND PIGS

LB677v1-Phar250wsg20kg-NZ0511